

Day 12 Plant Fossils Key

Fossil 1: *Dillhoffia cachensis*



About *Dillhoffia cachensis*:

- Approximate fossil time period: 49.5 MYA
- The fossil shows the part of the plant that creates a bud around a flower(sepal):a smaller, more complete one is on the upper left, and a larger, fragmented one. The **darkened areas are fruit**: the longer one is from *Dillhoffia*, and the others are stray fruits. **There are no petals present**.
- Climate around this time period saw a decline in global temperatures, going from a very warm to a cooler Earth.
- *Dillhoffia* is an extinct flowering plant from the **angiosperm** plant group.

The funnel-shaped sepals on a live tomato plant below are green. Petals on a flower are different colors. On this tomato plant they are yellow.



Fossil 2: *Annularia stellata*



About *Annularia stellata*:

- Approximate fossil time period: 310–280 MYA
- *Annularia stellata* leaves are arranged in whorls, similar to *Equisetum telmateia* (horsetails.)
- Three or more elongated leaves arranged in whorls radiate out from a central point, such as a stem.
- Leaves have one single vein.
- The climate was warm, with swampy land and dense forests.
- *Annularia stellata* is an extinct vascular horsetail plant.

This is a living horsetail plant.



Fossil 3 cycad (L) and fern (R)



About cycads and ferns:

- Approximate fossil time period: 165–00.5 MYA
- Both the cycad (**gymnosperm/vascular plant**) and fern (**vascular plant**) have simple pinnate-arranged leaves.
- **Cycad leaves are thin** and taper upward to a point. **Fern leaves appear more oval**, more cylindrical, with a tapered end and flat base.
- At the beginning of this time period, the climate was warm and wet, with vast tropical areas. By the end of this time period, the climate began heating up and reached levels much warmer than present day. Sea levels were high.

Living cycad



Living fern



Fossil 4: *Gingko biloba*



About *Gingko biloba*:

- Approximate fossil time period: 56–33.9 MYA
- *Gingko biloba* leaves are fan shaped. Veins are fork into pairs from the base of the leaf.
- The climate during this time period ranges from the warmest to the coldest. Ice begins to form at the polar regions around 34 MYA.
- *Gingko biloba* is a **gymnosperm**.

Living *Gingko biloba* leaves.



FIGURE CITE KEY

Figure 1

URL: https://commons.wikimedia.org/wiki/File:Dillhoffia_cachensis_SR_92-17-20_1.jpg

Attribution: Kevmin, CC BY-SA 3.0 <<https://creativecommons.org/licenses/by-sa/3.0/>>, via Wikimedia Commons

Tomato plant

https://commons.wikimedia.org/wiki/File:Tomato_fruit_and_flowers_at_day_52.jpg Dennis Brown, CC BY-SA 3.0 <<https://creativecommons.org/licenses/by-sa/3.0/>>, via Wikimedia Commons

Figure 2

URL: https://commons.wikimedia.org/wiki/File:Annularia_Stellata.jpg

Attribution: Woudloper, Public domain, via Wikimedia Commons

Horsetail plant

URL: <https://commons.wikimedia.org/wiki/File:Equisetopsida.jpg> Attribution: Rror, CC BY-SA 3.0

Figure 3

URL: https://commons.wikimedia.org/wiki/File:Early_Cretaceous_Plant_Fossils.jpg

Attribution: Georgialh, CC BY-SA 3.0 <<https://creativecommons.org/licenses/by-sa/3.0/>>, via Wikimedia Commons

Cyad

https://commons.wikimedia.org/wiki/File:Cycad_leaves_semicircle.jpg Attribution: Ian Alexander, CC BY-SA 3.0 <<https://creativecommons.org/licenses/by-sa/3.0/>>, via Wikimedia Commons

Fern

American royal fern (*Osmunda spectabilis*) has leaves similar to the fossil fern. URL: https://commons.wikimedia.org/wiki/File:Osmunda_spectabilis_close_up.jpg

Attribution: Ushakaron, CC0, via Wikimedia Commons

Figure 4

URL: https://commons.wikimedia.org/wiki/File:Ginkgo_biloba_Macabee_BC.jpg

Attribution: User:SNP(upload to en:wikipedia) ; User:tangopaso (transfer to Commons), CC BY-SA 3.0 <<https://creativecommons.org/licenses/by-sa/3.0/>>, via Wikimedia Commons

Gingko biloba

<https://commons.wikimedia.org/wiki/File:GinkgoLeaves.jpg> Attribution: Joe Schneid, Louisville, Kentucky, CC BY-SA 3.0 <<https://creativecommons.org/licenses/by-sa/3.0/>>, via Wikimedia Commons